



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Note on Linear Differential Equations with Constant Coefficients.

BY F. FRANKLIN.

1. In order to solve the differential equation

$$\alpha_0 \frac{d^m y}{dx^m} + \alpha_1 \frac{d^{m-1} y}{dx^{m-1}} + \dots + \alpha_m y \equiv f\left(\frac{d}{dx}\right) y = 0, \quad (1)$$

we may put $y = ve^{\alpha x}$, v being an undetermined function of x , as does M. Picard in his *Traité d'Analyse*, vol. III, p. 392. The result of this substitution in the first member of the equation is

$$e^{\alpha x} \left[f(\alpha) v + f'(\alpha) \frac{dv}{dx} + \dots + f^{(p)}(\alpha) \frac{d^p v}{dx^p} + \dots \right], \quad (2)$$

and hence, if α is a p -fold root of the algebraic equation $f(z) = 0$, it is plain that the given differential equation will be satisfied by putting

$$v = C_0 + C_1 x + \dots + C_{p-1} x^{p-1}, \quad (3)$$

where the C 's are arbitrary constants.

2. The solutions thus obtained, and corresponding to the various roots $\alpha_1, \alpha_2, \dots, \alpha_n$ of the equation $f(z) = 0$, appear to constitute a complete system of solutions of the differential equation, since they comprise m particular solutions of the type $x^q e^{\alpha x}$; but in order to prove that the system is really complete, it is necessary to show that there can exist no linear relation of the form

$$P_1 e^{\alpha_1 x} + P_2 e^{\alpha_2 x} + \dots + P_n e^{\alpha_n x} \equiv 0, \quad (4)$$

where the P 's are polynomials in x . M. Picard proves this in a manner sufficiently elementary, but somewhat artificial and involving considerable detail. The impossibility of the relation (4) may, however, be shown instantaneously and without any calculation.

3. We have only to observe that if v be a polynomial of the degree $p - 1$, so that $v, \frac{dv}{dx}, \dots$ are polynomials of descending degrees, the expression (2) does not vanish unless $f(\alpha) = f'(\alpha) = \dots = f^{(p-1)}(\alpha) = 0$, i. e. unless α is a p -fold root of $f(z) = 0$. Hence, in order that $y = Pe^{ax}$ be a solution of a differential equation (linear and with constant coefficients), P being a polynomial of degree $p - 1$, it is *necessary* as well as sufficient that α be a p -fold root of the auxiliary algebraic equation of that differential equation. Now $y = P_2 e^{a_2 x} + \dots + P_n e^{a_n x}$ is evidently a solution of a certain differential equation whose auxiliary algebraic equation does not possess the root α_1 ; but, by what has just been said, $y = P_1 e^{a_1 x}$ cannot be a solution of this last-named equation, and therefore cannot be identical with $-(P_2 e^{a_2 x} + \dots + P_n e^{a_n x})$, so that the relation (4) cannot exist. Q. E. D.